

# **Important Information for Healthcare Professionals to Remember About Deferasirox Abdi (deferasirox) Treatment**

This booklet provides detailed information on posology and monitoring of patients on deferasirox, to minimise key adverse effects including medication errors during treatment.

Adverse events should be reported. Reporting forms and information can be found at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard).

See deferasirox SmPC on [www.mhra.gov.uk](http://www.mhra.gov.uk) for further details.

Approved by MHRA:

## Table of contents

<b>1. What is deferasirox? .....</b>	<b>3</b>
<b>2. Formulation and method of administration .....</b>	<b>5</b>
<b>3. Dosing per indication – important differences to minimise the potential for medication errors .....</b>	<b>6</b>
3.1 Dosing for patients with non–transfusion-dependent thalassaemia (NTDT).....	6
3.2 Dosing for patients with chronic transfusional iron overload.....	7
<b>4. Safety and important monitoring requirements .....</b>	<b>9</b>
4.1 Unknown consequences of long-term use in paediatric patients .....	9
4.2 Dose-dependent rise in serum creatinine .....	9
4.3 Liver function test elevations <sup>1</sup> .....	12
4.4 Auditory (decreased hearing) <sup>1</sup> .....	12
4.5 Ocular disturbances (lens opacities) <sup>1</sup> .....	12
4.6 Overchelation in NTDT <sup>1</sup> .....	13
<b>5. Other monitoring recommendations &amp; actions<sup>1,2</sup>.....</b>	<b>14</b>
<b>6. Reporting suspected adverse reactions .....</b>	<b>16</b>

# 1. What is deferasirox?

## Licensed indications<sup>1</sup>

### *Chronic Transfusional Iron Overload*

Deferasirox is indicated for the treatment of chronic iron overload due to frequent blood transfusions ( $\geq 7$  ml/kg/month of packed red blood cells) in patients with  $\beta$ -thalassaemia major aged 6 years and older.

Deferasirox is also indicated for the treatment of chronic iron overload due to blood transfusions when deferoxamine therapy is contraindicated or inadequate in the following patient groups:

- In paediatric patients with  $\beta$ -thalassaemia major with iron overload due to frequent blood transfusions ( $\geq 7$  ml/kg/month of packed red blood cells) aged 2 to 5 years
- In adult and paediatric patients with  $\beta$ -thalassaemia major with iron overload due to infrequent blood transfusions ( $< 7$  ml/kg/month of packed red blood cells) aged 2 years and older
- In adult and paediatric patients with other anaemias aged 2 years and older

### *Non-Transfusion-Dependent Thalassaemia (NTDT)*

Deferasirox is also indicated for the treatment of chronic iron overload requiring chelation therapy when deferoxamine therapy is contraindicated or inadequate in patients with non-transfusion-dependent thalassaemia syndromes aged 10 years and older.

## **Mechanism of action<sup>1</sup>**

Deferasirox is an orally active chelator that is highly selective for iron (III). It is a tridentate ligand that binds iron with high affinity in a 2:1 ratio. Deferasirox promotes excretion of iron, primarily in the faeces. Deferasirox has low affinity for zinc and copper, and does not cause constant low serum levels of these metals.


## **Purpose of this booklet**

This booklet is for prescribers of deferasirox. It provides detailed information on posology and required monitoring of patients being treated with deferasirox, to minimise potential safety risks.

For full safety information, please refer to the deferasirox Summary of Product Characteristics.

## 2. Formulation and method of administration

Deferasirox is supplied as film-coated tablets which are available in three strengths<sup>1</sup>:



Tablets displayed are not actual size.

Deferasirox may be taken on an empty stomach or with a light meal.<sup>1</sup>

Deferasirox should be swallowed whole with some water. For patients who are unable to swallow whole tablets, deferasirox may be crushed and administered by sprinkling onto soft food (eg, yoghurt or apple sauce (apple puree)). The dose should be immediately and completely consumed, and not stored for future use.<sup>1</sup>

Deferasirox should be taken once a day, preferably at the same time each day.<sup>1</sup>

### 3. Dosing per indication – important differences to minimise the potential for medication errors

#### 3.1 Dosing for patients with non–transfusion-dependent thalassaemia (NTDT)

- Recommended initial dose of deferasirox: 7 mg/kg body weight/day<sup>1</sup>
- Doses >14 mg/kg/day are not recommended<sup>1</sup>
- Monitor your patients regularly to ensure proper treatment<sup>1</sup>

<b>Deferasirox: Starting dose and dose adjustment for patients with NTDT<sup>1</sup></b>			
<b>INITIATE</b> deferasirox <sup>a</sup>	<b>UP-TITRATE</b> to achieve target when necessary Monitor monthly	<b>DOWN-TITRATE</b> to avoid overchelation Monitor monthly	<b>STOP</b> chelation therapy once goal has been achieved
7 mg/kg/day	Increase in increments of 3.5 to 7 mg/kg/day up to a maximum dose of 14 mg/kg/day for adult patients and 7 mg/kg/day for paediatric patients <sup>a</sup>	Decrease dose to 7 mg/kg/day or less or closely monitor renal and hepatic function and serum ferritin levels	Re-treatment is not recommended for patients with NTDT
LIC <sup>b</sup> ≥5 mg Fe/g dw OR SF consistently >800 µg/l	LIC <sup>b</sup> ≥7 mg Fe/g dw OR SF consistently >2000 µg/l <sup>c</sup>	LIC <sup>b</sup> < 7 mg Fe/g dw OR SF consistently ≤2000 µg/l	<b>GOAL</b> LIC <sup>b</sup> <3 mg Fe/g dw OR SF consistently < 300 µg/l

dw, dry weight; LIC, liver iron concentration; NTDT, non–transfusion-dependent thalassaemia; SF, serum ferritin.

<sup>a</sup> Doses above 14 mg/kg/day are not recommended for patients with NTDT. In paediatric patients with NTDT, dosing should not exceed 7 mg/kg. In patients in whom LIC was not assessed, and SF is ≤2000 µg/l, dosing should not exceed 7 mg/kg.

<sup>b</sup>LIC is the preferred method of iron overload determination.

<sup>c</sup>In addition, a dose increase should only be considered if the patient is tolerating the medicinal product well.

### Paediatric NTDT patients<sup>1</sup>

In paediatric patients, dosing should not exceed 7 mg/kg/day. LIC should be monitored every 3 months when SF is  $\leq 800$   $\mu\text{g/l}$  in order to avoid overchelation.

WARNING: Data in children with NTDT are very limited. As a consequence, deferasirox therapy should be closely monitored to detect side effects and to follow iron burden in the paediatric population. In addition, before administering deferasirox to heavily iron-overloaded children with NTDT, the physician should be aware that the consequences of long-term exposure in such patients are currently not known.

### 3.2 Dosing for patients with chronic transfusional iron overload

- Recommended initial dose: 14 mg/kg body weight/day<sup>1</sup>
- Doses >28 mg/kg/day are not recommended<sup>1</sup>
- Monitor your patients regularly to ensure proper treatment<sup>1</sup>

<b>Deferasirox: Starting dose and dose adjustment for patients with transfusional iron overload<sup>1</sup></b>			
<b>INITIATE</b> Deferasirox therapy	<b>UP-TITRATE</b> to achieve goal when necessary Monitor monthly	<b>DOWN-TITRATE</b> to avoid overchelation Monitor monthly	<b>STOP</b> chelation therapy once goal has been achieved
<b>14 mg/kg body weight per day (recommended starting dose)</b> 20 U (~100 ml/kg) PRBCs or SF >1000 $\mu\text{g/l}$	Increase in increments of 3.5 to 7 mg/kg/day up to a maximum dose of 28 mg/kg/day	Decrease dose in steps of 3.5 to 7 mg/kg/day when SF = 500 to 1000 or closely monitor renal and hepatic function and serum ferritin levels	
<b>7 mg/kg body weight per day</b> <7 ml/kg/month of PRBCs (~ <2 units/month for an adult)	Increase in increments of 3.5 to 7 mg/kg/day up to a maximum dose of 28 mg/kg/day	—————	
<b>21 mg/kg body weight per day</b> >14 ml/kg/month of PRBCs	Increase in increments of 3.5 to 7 mg/kg/day up to a	Decrease dose in steps of 3.5 to 7 mg/kg/day when SF persistently <2500 $\mu\text{g/l}$ and showing a	

(~ >4 units/month for an adult)	maximum dose of 28 mg/kg/day	decreasing trend over time or closely monitor renal and hepatic function and serum ferritin levels	SF consistently <500 µg/l
<b>Patients already well managed on treatment with deferoxamine</b> Starting dose of deferasirox that is numerically one third that of the deferoxamine dose	Increase in increments of 3.5 to 7 mg/kg/day if dose is <14 mg/kg body weight per day and sufficient efficacy is not obtained	In patients treated with doses >21 mg/kg, decrease dose in steps of 3.5 to 7 mg/kg/day when SF persistently and <2500 µg/l showing a decreasing trend over time or closely monitor renal and hepatic function and serum ferritin levels	

PRBCs, packed red blood cells; SF, serum ferritin; U, units.

### Paediatric transfusional iron overload patients<sup>1</sup>

- The dosing recommendations for paediatric patients aged 2 to 17 years with transfusional iron overload are the same as for adult patients. Changes in weight of paediatric patients over time must be taken into account when calculating the dose
- In children with transfusional iron overload aged between 2 and 5 years, exposure is lower than in adults. This age group may therefore require higher doses than are necessary in adults. However, the initial dose should be the same as in adults, followed by individual titration
- It is recommended that serum ferritin be monitored every month to assess the patient's response to therapy and to minimise the risk of overchelation



## **4. Safety and important monitoring requirements**

### **4.1 Unknown consequences of long-term use in paediatric patients**

Data in children with NTDT are very limited. As a consequence, deferasirox therapy should be closely monitored to detect side effects and to follow iron burden in the paediatric population. In addition, before administering deferasirox to heavily iron-overloaded children with NTDT, the physician should be aware that the consequences of long-term exposure in such patients are currently not known.<sup>1</sup>

In paediatric patients with NTDT, dosing should not exceed 7 mg/kg/day. Liver iron concentration (LIC) should be monitored every 3 months when SF is  $\leq 800$   $\mu\text{g/l}$  in order to avoid overchelation.<sup>1</sup>

Body weight, height and sexual development testing should be conducted annually in paediatric patients.<sup>1</sup>

### **4.2 Dose-dependent rise in serum creatinine**

#### **Monitoring serum creatinine and creatinine clearance (CrCl)<sup>1</sup>**

Deferasirox may cause serious kidney problems, which can be fatal. Therefore, it is recommended that serum creatinine be assessed in duplicate before initiating therapy. Serum creatinine, CrCl (estimated with the Cockcroft–Gault or Modification of Diet in Renal Disease formula in adults and with the Schwartz formula in children), and/or plasma cystatin C levels should be monitored prior to therapy, weekly in the first month after initiation or modification of therapy with deferasirox, and monthly thereafter.

## Methods for estimating CrCl<sup>1</sup>

For your reference, here is a brief overview of methods to estimate CrCl in adults and children when prescribing deferasirox.

### **Adult**

Once a method has been selected, you should not change between or interchange formulas.

#### *Cockcroft–Gault formula<sup>2</sup>*

The Cockcroft–Gault formula employs creatinine measurements and the patient’s weight to predict CrCl.

The formula states CrCl in ml/min

$$\text{Estimated creatinine clearance in ml/minute} = \frac{(140 - \text{age}) \times \text{weight} \times \text{Constant}}{\text{Serum creatinine}}$$

**AGE = YEARS**

**WEIGHT = IDEAL BODY WEIGHT IN KG**

**SERUM CREATININE = MICROMOL/LITRE**

**CONSTANT = 1.23 FOR MEN; 1.04 FOR WOMEN**

#### *CKD-EPI equation<sup>3,4</sup>*

A general practice and public health perspective favours adoption of the CKD-EPI equation in North America, Europe, and Australia and using it as a comparator for new equations in all locations.

Glomerular filtration rate (GFR) =  $141 \times \min(\text{Scr}/\kappa, 1)^\alpha \times \max(\text{Scr}/\kappa, 1)^{-1.209} \times 0.993^{\text{Age}} \times 1.018$  [if female]  $\times 1.159$  [if black], where Scr is serum creatinine,  $\kappa$  is 0.7 for females and 0.9 for males,  $\alpha$  is  $-0.329$  for females and  $-0.411$  for males, min indicates the minimum of Scr/ $\kappa$  or 1, and max indicates the maximum of Scr/ $\kappa$  or 1.

### **Paediatric**

#### *Schwartz formula<sup>5</sup>*

$$\text{Creatinine clearance (ml/min)} = \frac{\text{constant}^b \times \text{height (cm)}}{\text{serum creatinine (mg/dl)}}$$

CKD-EPI, Chronic Kidney Disease Epidemiology Collaboration.

<sup>a</sup>If serum creatinine is provided in mmol/l instead of mg/dl, the constant should be 815 instead of 72.

<sup>b</sup>The constant is 0.55 in children and adolescent girls, or 0.70 in adolescent boys.

## Renal monitoring and actions<sup>1</sup>

	Serum creatinine		Creatinine clearance
<b>Monitoring before initiation of therapy</b>	Twice (2x)	and	Once (1x)
<b>Contraindicated</b>			<b>&lt;60 ml/min</b>
First month after start of therapy or dose modification	Weekly	and	Weekly
Thereafter	Monthly	and	Monthly
<b>Reduction of daily dose by 7 mg/kg/day, if following renal parameters are observed at two consecutive visits and cannot be attributed to other causes</b>			
Adult patients	>33% above pre-treatment average	and	Decreases <LLN (<90 ml/min)
Paediatric patients	>age appropriate ULN	and/or	Decreases <LLN (<90 ml/min)
<b>After dose reduction, interrupt treatment, if</b>			
Adult and paediatric	Remains >33% above pre-treatment average	and/or	Decreases <LLN (<90 ml/min)

Adapted from reference 1

LLN, lower limit of the normal range; ULN, upper limit of the normal range.

Treatment may be reinitiated depending on the individual clinical circumstances.

Dose reduction or interruption may be also considered if abnormalities occur in levels of markers of **renal tubular function** and/or as clinically indicated:

- Proteinuria (test should be performed prior to therapy and monthly thereafter)
- Glycosuria in patients without diabetes and low levels of serum potassium, phosphate, magnesium or urate, phosphaturia, aminoaciduria (monitor as needed).

Renal tubulopathy has been mainly reported in children and adolescents with  $\beta$ -thalassaemia treated with deferasirox.

Patients should be referred to a renal specialist, and further specialised investigations (such as renal biopsy) may be considered if the following occur despite dose reduction and interruption:

- Serum creatinine remains significantly elevated and
- Persistent abnormality in another marker of renal function (eg, proteinuria, Fanconi syndrome).

Consider hyperammonemic encephalopathy and early measurement of ammonia levels if patients develop unexplained changes in mental status while on deferasirox therapy, particularly in children.

### 4.3 Liver function test elevations<sup>1</sup>

Liver function test elevations have been observed in patients treated with deferasirox. Postmarketing cases of hepatic failure, sometimes fatal, have been reported in patients treated with deferasirox. Most reports of hepatic failure involved patients with significant morbidities including pre-existing liver cirrhosis. However, the role of deferasirox as a contributing or aggravating factor cannot be excluded.

If there is a persistent and progressive increase in serum transaminase levels that cannot be attributed to other causes, deferasirox should be interrupted. Once the cause of the liver function test abnormalities has been clarified or after return to normal levels, cautious reinitiation of treatment at a lower dose followed by gradual dose escalation may be considered.

#### Monitoring requirements for liver function tests

Monitor	Frequency <sup>1</sup>
Serum transaminases Bilirubin Alkaline phosphatase	Serum transaminases, bilirubin and alkaline phosphatase should be checked <b>prior to therapy, every 2 weeks during the first month and monthly thereafter</b>

Consider hyperammonemic encephalopathy and early measurement of ammonia levels if patients develop unexplained changes in mental status while on deferasirox therapy, particularly in children.

### 4.4 Auditory (decreased hearing)<sup>1</sup>

Auditory (decreased hearing) disturbances have been reported in patients treated with deferasirox, however they are uncommon.

Auditory testing is recommended before the start of treatment and at regular intervals thereafter (every 12 months).

If disturbances are noted during the treatment, dose reduction or interruption may be considered.

Monitoring	Frequency <sup>1</sup>	Action
Auditory	Auditory monitoring recommended prior to therapy and yearly thereafter	If disturbances in hearing during treatment, consider dose reduction or interruption

### 4.5 Ocular disturbances (lens opacities)<sup>1</sup>

Ocular disturbances (lens opacities) have been reported in patients treated with deferasirox, however they are uncommon.

Ophthalmic testing (including fundoscopy) is recommended before the start of treatment and at regular intervals thereafter (every 12 months).

If disturbances are noted during the treatment, dose reduction or interruption may be considered.

<b>Monitoring</b>	<b>Frequency<sup>1</sup></b>	<b>Action</b>
Ophthalmic (including funduscopy)	Ophthalmic monitoring recommended prior to therapy and yearly thereafter	If disturbances in vision during treatment, consider dose reduction or interruption

## 4.6 Overchelation in NTDT<sup>1</sup>

Chelation therapy should only be initiated when there is evidence of iron overload (LIC  $\geq$ 5 mg Fe/g dry weight [dw] or serum ferritin consistently  $>$ 800  $\mu$ g/l). LIC is the preferred method of iron overload determination and should be used wherever available. Caution should be taken during chelation therapy to minimise the risk of overchelation in all patients.

In paediatric patients with NTDT, dosing should not exceed 7 mg/kg/day. In these patients, closer monitoring of LIC and serum ferritin is essential to avoid overchelation. In addition to monthly serum ferritin assessments, LIC should be monitored every 3 months when serum ferritin is  $\leq$ 800  $\mu$ g/l.<sup>1</sup>

<b>Monitor</b>	<b>Frequency<sup>1</sup></b>	<b>Action</b>
Serum ferritin (SF)	Prior to therapy and monthly thereafter	If SF $<$ 300 $\mu$ g/l, interrupt treatment
Liver iron concentration (LIC)	All patients: Prior to therapy Paediatric patients only: Every 3 months if SF is $\leq$ 800 $\mu$ g/l	If LIC $<$ 3 mg Fe/g dw, interrupt treatment

## 5. Other monitoring recommendations & actions<sup>1</sup>

Please refer to table below for treatment interruption conditions.

Consideration	Treatment interruption conditions
SF	Consistently <500 µg/l (in transfusional iron overload) or <300 µg/l (in NTDT syndromes)
Serum creatinine	Adult and paediatric: after dose reduction, when serum creatinine remains >33% above baseline and/or CrCl <LLN (90 ml/min) – also refer patient to renal specialist and consider biopsy
Proteinuria	Persistent abnormality – also refer patient to renal specialist and consider biopsy
Tubular markers	Abnormalities in levels of tubular markers and/or if clinically indicated – also refer patient to renal specialist and consider biopsy (also consider dose reduction)
Serum transaminases (ALT and AST)	Persistent and progressive increase in liver enzyme levels
Metabolic acidosis	Development of metabolic acidosis
SJS, TEN, or any other severe skin reaction (eg, DRESS)	Suspicion of any Severe Cutaneous Adverse Reaction (SCAR): discontinue immediately and do not reintroduce
Hypersensitivity reactions (eg. anaphylaxis, angioedema)	Occurrence of reaction: discontinue and institute appropriate medical intervention. Do not reintroduce in patients who have experienced a hypersensitivity reaction due to the risk of anaphylactic shock
Vision and hearing	Disturbances during the treatment (also consider dose reduction)
Unexplained cytopenia	Development of unexplained cytopenia

ALT, alanine aminotransferase; AST, aspartate aminotransferase; CrCl, creatinine clearance; DRESS, drug reaction with eosinophilia and systemic symptoms; LLN, lower limit of the normal range; NTDT, non-transfusion-dependent thalassaemia; SF, serum ferritin; SJS, Stevens–Johnson syndrome; TEN, toxic epidermal necrolysis.

Please refer to the table below for appropriate monitoring and disease markers.

	<b>Baseline</b>	<b>In the first month after initiation of deferasirox or after dose modification</b>	<b>Monthly</b>	<b>Every 3 months</b>	<b>Yearly</b>
SF	✓		✓		
LIC <sup>a</sup>	✓			✓ (for paediatric patients with NTDT only, if SF is ≤800 µg/l)	
Serum creatinine	<b>2x</b>	<b>Weekly</b> (Should also be tested weekly in the first month after dose modification)	✓		
Creatinine clearance and/or plasma cystatin C	✓	<b>Weekly</b> (Should also be tested weekly in the first month after dose modification)	✓		
Proteinuria	✓		✓		
Serum transaminases, bilirubin, alkaline phosphatase	✓	<b>Every 2 weeks</b>	✓		
Body weight, height, and sexual development	✓				✓ <sup>b</sup>
Auditory/ophthalmic testing (including fundoscopy)	✓				✓

LIC, liver iron concentration; SF, serum ferritin.

<sup>a</sup>For patients with NTDT, LIC is the preferred method of iron overload determination and should be used wherever available. Caution should be taken during chelation therapy to minimise the risk of overchelation in all patients.

<sup>b</sup> Paediatric patients only.

The results of the tests for serum creatinine, CrCl, plasma cystatin C, proteinuria, serum ferritin, liver transaminases, bilirubin, and alkaline phosphatase should be recorded and regularly assessed for trends. The results should also be noted in the patient's charts, along with pre-treatment baseline levels for all tests.<sup>1</sup>

## **6. Reporting suspected adverse reactions**

Please report suspected adverse drug reactions (ADRs) to the MHRA through the Yellow Card Scheme. You can report via the Yellow Card Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) The free Yellow Card app is available from the Apple App Store or Google Play Store. Some clinical IT systems (EMIS/SystemOne/Vision/MiDatabank) for healthcare professionals.

Alternatively, you can report a suspected side effect to the Yellow Card Scheme by calling 0800 731 6789 for free, Monday to Friday between 9am and 5pm. You can leave a message outside of these hours.

When reporting please provide as much information as possible. By reporting side effects, you can help provide more information on the safety of this medicine.

Adverse events should be reported. Reporting forms and information can be found at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard).



**References:**

1. Deferasirox Summary of Product Characteristics
2. Cockcroft DW, Gault MH. *Nephron*. 1976;16(1): 31–41.
3. Earley A, Miskulin D, Lamb EJ, Levey AS, Uhlig K. *Ann Intern Med*. 2012;156(11):785–795.
4. Levey AS, Stevens LA, Schmid CH, et al.; for the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI). *Ann Intern Med*. 2009;150(9):604–612.
5. Schwartz GJ, Brion LP, Spitzer A. *Pediatr Clin North Am*. 1987;34(3):571–590.